

An original constraint on the Hubble constant : $h > 0.74$

A. Barrau^{1*}, A. Gorecki¹ and J. Grain²

¹*Laboratoire de Physique Subatomique et de Cosmologie, Université Joseph Fourier, CNRS/IN2P3, INPG
53, avenue des Martyrs, 38026 Grenoble cedex, France*

²*Laboratoire AstroParticule et Cosmologie, Université Paris 7, CNRS/IN2P3
10, rue Alice Domon et Léonie Duquet, 75205 Paris cedex 13, France*

Accepted xxxx Received xxxx; in original form xxxx

ABSTRACT

The Hubble parameter H_0 is still not very well measured. Although the Hubble Key Project, Chandra and WMAP gave good estimates, the uncertainties remain quite large. In this brief report, we suggest an original and independent method to derive a lower limit on H_0 using the absorption of very high energy gamma-rays by the cosmic infrared background. With conservative a hypothesis, we obtain $H_0 > 74 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ at the 68% confidence level, which favors the upper end of the intervals allowed by dedicated experiments.

Key words: cosmological parameters – diffuse radiation.

INTRODUCTION

Gamma-rays in the $10^{11} - 10^{13}$ eV range are observed from distant Active Galactic Nuclei (AGN). Those very high-energy photons will be absorbed by the Cosmic Infrared Background (CIB) due to electron-positron pair production. In the higher energy range (typically around 10-20 TeV), the absorption is expected to be so high that the difference between the source spectrum and the measured spectrum becomes very large. As the CIB spectral distribution is now quite correctly known (at least with lower limits) it is possible to compute the intrinsic spectrum of a given AGN as a function of the integrated CIB density crossed along the line-of-sight through an unfolding procedure. The absorption being proportional to e^{1/H_0} , it is then possible to derive a sensitive lower limit on the Hubble parameter so as to exclude unfolded spectra which would be explicitly unphysical. The idea is very simple : the smaller the Hubble constant, the larger the distance to the source, the larger the number of CIB photons crossed, and the higher the unfolding coefficients. As the unfolded (*i.e.* intrinsic) spectrum shape will inevitably become unacceptable for low enough Hubble parameters, this approach allows to derive a lower limit on H_0 . The obtained bound is relevant when compared with current estimates, although the main aim of this brief article is to open this field and outline the scheme of the method. In the first section, the latest measurements of the CIB density are reviewed, including some recently revised analysis. In the second section, the gamma-ray data relevant for this study are presented together with the associated uncertainties. In the third section, the physical

processes involved in the gamma-ray absorption mechanism are explained and the basic features of the proposed method are underlined. The fourth section deals with the details of the unfolding procedure and with the physical criteria used to reject "unphysical" unfolded spectra. In the fifth section, the Monte-Carlo method used to derive statistically meaningful bounds is described and the results are given. Finally, several developments that could be expected in the future are outlined in the last section.

1 THE COSMIC INFRARED BACKGROUND DENSITY

Our understanding of the early epochs of galaxies has recently increased thanks to the observational evidences provided by UV/Visible/Near-IR, far-IR and submillimeter surveys of high-redshift objects. In a consistent scenario, galaxy formation and evolution can also be constrained by the background radiation which is produced by the line-of-sight accumulation of all extragalactic sources. The Cosmic Infrared Background (CIB) is basically the relic emission at wavelengths between a few microns and millimeters of the formation and evolution of galaxies of all type and star-forming systems (see, *e.g.*, Puget *et al.* (1996), Hauser *et al.* (1998), Lagache *et al.* (1999), Gispert *et al.* (2000), Hauser & Dwek (2001) and Kashlinsky (2005)). The near-IR CIB arises mainly from the stellar component of galaxies and probes their evolution at early times. The mid- and far-IR CIB originates from dusty galaxies reprocessing stellar light and other energetic output.

* E-mail: Aurelien.Barrau@cern.ch

Observationally, the CIB is difficult to distinguish from the generally brighter foregrounds contributed by the local matter within the solar system, the stars and the interstellar medium of the Galaxy. However, the situation has dramatically improved in the last decade. In this study, we have used the most accurate and "up-to-date" estimates of the CIB density, as described, *e.g.*, in Dole *et al.* (2006) with linear interpolations (in log-log scale) between the measurements. The doubtful measurements have deliberately been ignored. The number of points is now large enough to make this approach quite accurate. The measurements used in our analysis are the following:

- $0.3 \mu\text{m}$: $10 \pm 5 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from Bernstein *et al.* (2002) corrected by Matilla (2003). The estimate is based on the Hubble Space Telescope (HST) broadband CCD photometry and the 2.5 m du Pont telescope from Lac Campanas Observatory (LCO).
- $1.25 \mu\text{m}$: $28 \pm 15 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from Wright (2001) and Cambresy *et al.* (2001). The estimate relies on the Diffuse Infrared Background Experiment (DIRBE) and 2 Micron All Sky Survey (2MASS) data.
- $2.2 \mu\text{m}$: $22.4 \pm 6 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from Gorjian *et al.* (2000) using DIRBE and *Lick* data.
- $3.3 \mu\text{m}$: $11.1 \pm 3.3 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from the same analysis.
- $15 \mu\text{m}$: $3.3 \pm 1.3 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from Elbaz *et al.* (1998). This lower limit is based on galaxy counts with the Infrared Space Observatory Camera (ISOCAM).
- $24 \mu\text{m}$: $2.7 \pm 1.1 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from Papovich *et al.* (2004). The limit comes from galaxy counts with the Multi-band Imaging Photometer for *Spitzer* (MIPS).
- $70 \mu\text{m}$: $7.1 \pm 1.0 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from Dole *et al.* (2006), based on MIPS data with more than 19000 $24 \mu\text{m}$ sources stacked with $S_{24} > 60 \mu\text{Jy}$.
- $160 \mu\text{m}$: $13.4 \pm 1.7 \text{ nW.m}^{-2}.\text{sr}^{-1}$, from the same analysis.

It is important to underline that most of those estimates are *lower limits*. As the constraints derived on the Hubble parameter in the next sections would have only been weakened by a lower CIB density, this makes the following analysis conservative. The *Spitzer* observatory data (Werner *et al.* (2004)) allowed for deep and wide area surveys, in particular at 24, 70 and $160 \mu\text{m}$ using the MIPS (Rieke *et al.* (2004)) which is a keypoint for this analysis. Those points, used here with the refined analysis of Dole *et al.* (2006), improving over Dole *et al.* (2004), are very relevant as they (at least two of them) lie within the energy range where the interactions with gamma-rays are large. In the following, we therefore consider the CIB density as known (within the uncertainties) and use it as an "absorbing material" for the gamma-rays. The higher the distance to the source (*i.e.* the lower the Hubble parameter), the higher the absorption. As the transmitted flux depends exponentially on the depth of the absorber, the approach is very sensitive.

Fig. 1 displays the data points (filled circles) used for this analysis together with lower limits (triangles) and estimates not taken into account as being affected by higher uncertainties (open circles). The Monte-Carlo method de-




Figure 1. Cosmic infrared background energy density as a function of the wavelength. The filled circles are the data selected for this analysis. Open circles are tentative detections ignored in this article and triangles are lower limits. The intervals labeled "4 TeV sensitivity" and "17 TeV sensitivity" correspond to 90% of the interactions of, respectively, 4 TeV and 17 TeV gamma-rays taking into account the cross section convolved with the CIB spectrum. The shaded region corresponds to the 1σ allowed values of the CIB density, as included in the Monte-Carlo simulation.

scribed in the next sections samples CIB realizations within the shaded region.

2 ACTIVE GALACTIC NUCLEUS MRK 501

It has been suggested long ago that observations of the TeV spectrum of extragalactic sources can be a powerful tool to constrain the CIB spectrum, especially around $10 \mu\text{m}$, where the Extragalactic Background (EB) constraints were quite weak. As a matter of fact, TeV gamma rays propagating in the intergalactic medium undergo absorption through electron-positron pair production on CIB photons. Using sources Mrk 421 and Mrk 501, meaningful upper limits have been established in different papers. More recently, interesting constraints were obtained by Aharonian *et al.* (2006). They are based on the observations performed with the very effective new network of telescopes H.E.S.S. (Aharonian *et al.* (2005a)). In this article, we reverse the argument and use reliable estimates of the CIB density to obtain a new constraint on the Hubble parameter. The idea has already been pointed out by Salamon *et al.* (1994). However, Mrk 501 data were not available at that time and no result could be derived. Furthermore, the method suggested by Salamon *et al.* is far less conservative as it assumes that any rolloff in the AGN spectrum should be due to the CIB absorption. This is a very strong hypothesis as many different effects could (and, even, should) lead to a rolloff intrinsic to the source. This could, among other processes, be due to self-absorption within the AGN, to the Klein-Nishina effect or simply to the limiting energy available for Fermi acceleration to take place in a finite volume over a finite amount of time. The bound derived in this article does not rely on this "CIB rolloff" assumption and is therefore much more robust. But it can only lead to

a lower limit and not to a measurement.

The TeV source Mrk 501 is the second closest X-ray selected BL Lac object after Mrk 421, with a redshift $z = 0.034$. During the 1997 outburst which lasted several months, Mrk 501 was observed intensively in X-rays (BeppoSAX: Pian *et al.* (1998), RXTE: Lamer & Wagner (1998)) and TeV γ -rays (Whipple: Catanese *et al.* (1997), Samuelson *et al.* (1998); HEGRA: Aharonian *et al.* (1997), Telescope Array: Hayashida *et al.* (1998); CAT: Djannati-Ataï *et al.* (1999)). The exceptional 1997 April 16 flare was observed by BeppoSAX and low-energy threshold (~ 300 GeV) Whipple and CAT atmospheric Cherenkov telescopes, allowing the derivation of the energy spectrum with a good accuracy in a broad dynamical range. These unique data initiated interesting efforts to set meaningful upper limits on the CIB flux (see *e.g.* Biller *et al.* (1998), Stanev & Franceschini (1998), Stecker & De Jager (1998), Aharonian *et al.* (1999b), Coppi & Aharonian (1999), Konopelko *et al.* (1999), Guy *et al.* (2000), Renault *et al.* (2001)). More recently, other AGNs have been detected, especially with the H.E.S.S. network (see, *e.g.*, Aharonian *et al.* (2005b) for a review). As far as our study is concerned, a promising opportunity is the measurement of the spectrum of the distant blazar 1ES1101-232 (Aharonian *et al.* (2007)). With a redshift $z = 0.186$, this is a very good candidate to maximize the interaction with the CIB. The spectrum has been measured by H.E.S.S. up to $\sim 2 - 3$ TeV. In fact, the optical depth for gamma-rays is roughly the same at this energy for $z = 0.186$ as at $\sim 15 - 20$ TeV for the lower Mrk 501 redshift $z = 0.034$. The analysis hereafter described has also been performed for both Mrk 501 and 1ES1101-232. However, as the results are less stringent for the latter, we will mainly focus on Mrk 501. Furthermore, the very wide range of measured energies (between ~ 400 GeV and ~ 20 TeV) for Mrk 501 allows for a much more reliable analysis. In this article, CAT and HEGRA data are used from 400 GeV to 21 TeV (Aharonian *et al.* (1999b) and Guy *et al.* (2000)). The higher energy points, *i.e.* the more relevant ones for this analysis, come from a time averaging of the HEGRA spectra based on more than 38000 detected γ -rays. This very high number of data points combined with the 20% energy resolution of the instrument resulted in a very accurate measurement of the spectrum above 10 TeV which remains unequaled.

3 INTERACTIONS OF GAMMA-RAYS WITH THE INFRARED BACKGROUND

The influence of low energy photons in the Universe on the propagation of Very High Energy (VHE) gamma-rays was first pointed out by Nikishov (1962). An original way of using ground-based TeV observations of distant sources to probe the CIB was given by Stecker *et al.* (1992). The fundamental idea is to look for absorption in the intrinsic spectrum as a result of electron-positron pair production by photon collisions $\gamma_{TeV} + \gamma_{CIB} \rightarrow e^+ + e^-$. In such an interaction between a gamma-ray of energy $(1+z)E$ and an infrared photon of energy $(1+z)\epsilon$, with E and ϵ the observed energies at $z = 0$, the pair production threshold is $E\epsilon(1+z)^2(1-\cos\theta) > 2(mc^2)^2$ where θ is the

angle between photons and m the rest mass of the electron. The cross-section can be written as (Heitler (1960)): $\sigma = \zeta(1-\beta^2)(2\beta(\beta^2-2) + (3-\beta^4)\ln(\frac{1+\beta}{1-\beta}))\text{cm}^2$ with $\beta = \left(1 - 2(mc^2)^2/(E\epsilon(1-\cos\theta))(1+z)^2\right)^{1/2}$ and $\zeta = 1.25 \times 10^{-25}$. If the infrared photons have a density number $n(\epsilon)d\epsilon\text{cm}^{-3}$, the corresponding optical depth for attenuation is

$$\tau(E) \approx \frac{cz_s}{H_0} \int_{-1}^1 d(\cos\theta) \frac{1-\cos\theta}{2} \int_{\epsilon_t}^{\infty} d\epsilon n(\epsilon)\sigma(E, \epsilon, \theta)$$

where $\epsilon_t = 2(mc^2)^2/(E(1-\cos\theta)(1+z)^2)$, z_s is the redshift of the source and c the speed of light. This expression is strictly valid for an Einstein-de Sitter universe but remains an accurate approximation in the Λ CDM paradigm for low redshifts. The detected flux is then attenuated by a factor of $e^{-\tau(E)}$. The CIB energy distribution is assumed to be independent of z as the γ -ray source redshift is very low (0.034). The maximum cross-section is reached for an infrared photon wavelength of $\lambda_{CIB} \approx \lambda_c \frac{E}{2mc^2}$ where $\lambda_c = h/(mc)$ is the Compton wavelength of the electron. Therefore, γ -photons with energy between a few TeV and 20 TeV "see" CIB photons with wavelengths between 3.5 and 100 μm .

To allow for an intuitive understanding of the exponential absorption (not only as a function of the quantity of the absorbing material, as it should be, but also as a function of the energy of the gamma-rays), one can assume that the CIB is roughly constant, *i.e.* $\epsilon^2 n(\epsilon) \approx cte$. The optical depth for an energy kE can then be written as:

$$\tau(kE) \propto \int_{-1}^1 d(\cos\theta) \frac{1-\cos\theta}{2} \int_{\frac{2(mc^2)^2}{kE(1+z)^2(1-\cos\theta)}}^{\infty} d\epsilon \frac{\sigma(kE\epsilon, \theta)}{\epsilon^2}.$$

By a change of variable $\epsilon \rightarrow k\epsilon$, it is straightforward to see that $\tau(kE) = k \times \tau(E)$. As $\Phi = \Phi_0 \times e^{-\tau(E)}$, this explains why a high energy gamma-ray will be exponentially more absorbed than a low energy one.

4 UNFOLDING METHOD AND PHYSICAL CRITERIA

To turn the absorption corrected spectra into a lower limit on the Hubble parameter, the intrinsic spectral energy density of Mrk 501 is assumed to be concave (*i.e.* with a negative second derivative) in the multi-TeV region. This quite common approach has been developed in Renault *et al.* (2001) and Guy *et al.* (2000). The conservative hypothesis is based on the fact that no natural physical process can re-inject energy above the Inverse-Compton bump maximum. Both Klein-Nishina effect and auto-absorption within the source would only cause the flux to decrease more and more rapidly as a function of energy. This is obviously true for leptonic models (a good review of theoretical perspectives, where each model produces a concave spectrum, can be found in Saugé (2004)) but also for more complex hadronic models where this concave shape is also expected, either because of proton-initiated-cascade (often modeled as a broken power-law around 3 TeV), or due to the inclusion of μ -synchrotron radiation (Mannheim,

private communication). Even in the extreme case where proton synchrotron radiation is at the origin of the TeV bump (Aharonian (2000)), the shape of the emission remains concave. Looking at the absorption corrected spectra, it cannot be totally excluded that the maximum of the so-called Inverse-Compton peak is not yet reached at 20 TeV. Physical parameters (essentially magnetic field and Doppler factor) required to produce such a spectral energy distribution maximum above 17 TeV are substantially disfavored but, even in this case, the $\nu F(\nu)$ shape should remain concave. This latter point is demonstrated by BeppoSAX 1-100 keV measurements (Pian *et al.* (1998)) showing a clearly concave spectrum before the synchrotron bump maximum (around 100 keV) which is supposed to be mimicked by the TeV-spectrum before the so-called Inverse-Compton bump maximum. This hypothesis is reinforced by the fact that the sub-TeV slope, which is independent of the CIB density beyond $3.5 \mu\text{m}$, effectively reflects the X-ray slope in the keV range. This behavior (together with the correlated variability reported *e.g.* by Aharonian *et al.* (1999) and Djannati-Ataï *et al.* (1999)) indicates that the same population of particles is at the origin of both X-ray and γ -ray emissions, whatever this population is. In particular, the self-synchro-Compton model fits satisfactorily the absorption corrected data (Guy *et al.* (2000)).

Furthermore, in addition to those "theoretically motivated" arguments, observations show that the spectrum should indeed be concave. This can be checked by considering AGNs with an IC bump seen at lower energies and therefore unaffected by the CIB absorption. This can easily be seen for 3C273 with the average spectrum compiled from 30 years of observations as reported in Türler *et al.* (1999). This is also true for Cen A (see, *e.g.*, Chiaberge *et al.* (2001)) which is the nearest radiogalaxy ($z=0.0018$) and one of the best studied. Although with less significance this can even be checked for M 87 (which is not a blazar), as shown in Lenain *et al.* (2008). The FSRQ PKS 0521-36 (see, *e.g.* the summary presented in Lenain *et al.* (2008), mostly from Giommi *et al.* (2002) and the NED), which is known for oscillating between a Seyfert-like and a BL Lac state, also exhibits a fully concave spectrum. This is also true for the blazar 3C 454.3, as measured during the 2007 July flare (Ghisellini *et al.* (2007)). During flares, good quality data reinforcing the concavity assumption are also available for PKS 2155-304 (Aharonian *et al.* (2007)) and for 3C 279 (see, *e.g.*, the summary plot in Böttcher *et al.* (2007)).

It should be stressed that requiring the spectrum to be concave is "more demanding" than simply requiring the end of the TeV spectrum not to exhibit a too hard spectral index, as often used to derive limits on the CIB. Our approach evades the argument of Stecker *et al.* (2007) questioning the results of Aharonian *et al.* (2006).

In order to quantify the concavity of the absorption corrected spectrum, a parabolic fit is performed from 6 to 21 TeV in the plane $(\log(\nu), \log(\nu F\nu))$. This function, which is simple, is chosen for its constant second derivative $a = d^2(\log(\nu F\nu))/d^2(\log(\nu))$ which avoids the choice of a particular test-energy and fits satisfactorily the data. Over such a small interval it is obviously meaningful to




Figure 2. Unfolded gamma-ray spectra and associated fits for (from bottom to top) : no CIB (equivalent to $H_0 \rightarrow \infty$), $H_0 = 90 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ and $H_0 = 50 \text{ km.s}^{-1}.\text{Mpc}^{-1}$.

assume the spectrum to be locally parabolic. The previous physical constraint on the concavity of the TeV spectrum simply reads as $a < 0$. As explained in the next section, the uncertainties are carefully taken into account. The parameter a is computed for different values of the Hubble parameter and the corresponding value is rejected if a is positive within the associated errors. Fig. 2 shows the result of the parabolic fit in several cases, superimposed with the experimental points. The fit under-estimates the energy density of the hardest photons, ensuring that the local second derivative is under-estimated, thus making the test conservative.

5 RESULTS ON THE HUBBLE PARAMETER

To derive statistically meaningful constraints on the Hubble parameter, errors on both the CIB and the measured TeV spectrum have been taken into account. As far as the gamma-ray spectrum is concerned, we have not only considered the statistical uncertainties but also the systematic errors, as evaluated by the HEGRA team (Aharonian *et al.* (1999b)). The errors on the CIB density are not as easy to take into account as those on the AGN spectrum as their propagation is far from trivial. For this reason, we have performed a full Monte-Carlo simulation. Each of the eight considered CIB points between 0.3 and $160 \mu\text{m}$ is randomly generated according to its central value and uncertainty, as displayed by the shaded region of Fig.1. Linear interpolations are then performed in the log-log plane to obtain a CIB realization. For each CIB realization, the unfolded spectrum is computed (*i.e.* the optical depth is numerically evaluated for the considered energies with the formula given in section 3) and fitted with a parabola between 6 and 21 TeV. The error Δa on the second derivative a of

the parabola (due to uncertainties in the measurement of the gamma-ray spectrum) is computed according to three different statistical methods iterated up to the convergence point. The value of the Hubble parameter is rejected at $x\sigma$ if $a - x\Delta a > 0$. The procedure is finally repeated to generate many (typically ~ 1000) realizations of the CIB. Confidence levels can be computed in this way by estimating the percentage of rejected realizations for each tested value of the Hubble parameter. It has been checked that our results are marginally dependent upon the number of points used in the fit and that the χ^2 per degree of freedom is, in each case, low enough to ensure a satisfactory fit quality.

At the 68% confidence level, the lower limit derived on H_0 , taking into account all the uncertainties is: $H_0 > 74 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. This is the main result of this article. Although it should be underlined that even if a 1σ limit remains rather weak on a statistical basis, this bound is meaningful –at this confidence level– when compared with other results. The history of the measured value of the Hubble parameter is very chaotic and different (incompatible) values have been taken as granted at different times. As reminded in the recent review article “the Hubble constant” by Jackson (2007), the current estimates range between 60 and $75 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. In this context, our lower limit $H_0 > 74 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ is unquestionably relevant.

As far as Cepheids are concerned, the main measurements are $H_0 = 73 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ (Riess *et al.* (2005)), with statistical errors of $4 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ and systematic errors of $5 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ on the one hand and $H_0 = 62.3 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ (Sandage *et al.* (2006)), with statistical errors of $1.3 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ and systematic errors of $5 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ on the other hand. Our results clearly disfavor the Sandage estimate, even within its quoted uncertainty. As far as gravitational lenses are concerned, a sophisticated meta-analysis has recently been performed (Oguri *et al.* (2007)) using Monte-Carlo methods to account for quantities such as the presence of clusters around the main lens galaxy and the variation in profile slopes. The result obtained is $H_0 = 68 \pm 6 \pm 8 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. Meanwhile, an approach modelling simultaneously 10 of the 18 time-delay lenses with non-parametric models, has led (Saha *et al.* (2006)) to $H_0 = 72 \pm 12 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. Our results allow a significant 1σ narrowing of those uncertainty intervals. As far as the Sunyaev-Zeldovich effect is concerned, recent measurements (see, *e.g.* Jones *et al.* (2005)) are compatible with those estimates but the errors are too large to allow for any improvement.

The most reliable measures available to date are probably those reported in the “final results from the Hubble Space Telescope Key Project” (Freedman *et al.* (2001)). They concluded that the full analysis of all data gives $H_0 = 72 \pm 8 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. Combining this HST allowed interval with our lower limit increases the mean value to $H_0 \approx 76 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. However, the statistical significance of this estimate should be taken with care as the accurate distribution is not known. Nevertheless, our result $H_0 > 74 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ clearly favors the upper end of the HST interval and substantially reduces the allowed




Figure 3. Lower limit at the 90% confidence level on the Hubble constant as a function of the rescaling factor of the CIB density. A factor of x means that the errors are assumed x times smaller than currently estimated. Each point is obtained by analysing 1000 Monte-Carlo realizations of the CIB.

parameter space. A less constraining, although interesting, result was obtained by Chandra (Bonamente *et al.* (2006)) leading to $H_0 = 77 \pm 12 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ with a larger 15 % uncertainty. Finally, WMAP has published in the five-years results (Komatsu *et al.* (2008)) an impressive estimate at $H_0 = 72 \pm 2.6 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. It should, however, be strongly emphasized that WMAP results rely on many different priors and do not allow for an independent determination of H_0 due to its degeneracy with the total curvature of the Universe. For example, every decrease of $20 \text{ km.s}^{-1}.\text{Mpc}^{-1}$ in H_0 increases the total density of the Universe by 0.1 in units of the closure density. The WMAP data by themselves, without any further assumptions or extra hypothesis, do not supply a significant constraint on H_0 .

If the statistical significance is to be improved to 90%, the lower limit derived in this work is significantly reduced to $H_0 > 56 \text{ km.s}^{-1}.\text{Mpc}^{-1}$. This is, of course, due to large uncertainties on the CIB density. This broadening is somehow analogous to what happens for other approaches to measure H_0 . For example, Tegmark *et al.* (2006) have shown that when relaxing the constraint of the equation of state of the dark energy, the 90% confidence level estimate using WMAP data is $H_0 \in [61 - 84] \text{ km.s}^{-1}.\text{Mpc}^{-1}$.

CONCLUSION AND PROSPECTS

The lower limit on H_0 derived in this letter is totally *independent* of the other available measurements. It is conservative, both due to the method which tends to under-estimate the high energy flux density and to the weak hypothesis regarding the spectrum concavity. Internal effects, such as Klein-Nishina cutoff and self-absorption, lead to very sharp multi-TeV intrinsic spectra. Furthermore, and most importantly, the CIB density is underestimated as only lower limits are used. Nevertheless, the errors on those lower limits have been exhaustively taken into account thanks to a Monte-Carlo simulation. Our result

allows a substantial 1σ narrowing of previously obtained uncertainty intervals, favoring the higher possible values of H_0 . It also closes the window on exotic "low Hubble constant" scenarios (Nugent *et al.* (1995)) and on some alternatives to the cosmological concordance model (see, *e.g.*, Blanchard *et al.* (2003) where $h \approx 0.46$ is assumed.)

Clearly, this work aims at providing a first hint in this direction and a first order of magnitude of what can be obtained by this method. Although the statistical significance of our results has been computed with care, several improvements could be expected in the future. First, other AGNs will probably be detected by H.E.S.S. (Aharonian *et al.* (2005a)), MAGIC (Bastieri (2006)) or VERITAS (Krennrich (2007)) and should be included in the analysis. GLAST (Latronico *et al.* (2007)) data, at lower energies, could even be useful. This is very promising to account for more sources at different redshifts. However, the current situation for known AGNs is not expected to be improved significantly for a simple reason: the most important part of the spectrum used to constrain the Hubble parameter is the high-energy tail and larger telescopes will not improve the sensitivity in this range (just because the effective area is determined by the area of the shower and not by the area of the mirror). Then, improvements can be expected from measurements of the CIB. Fig 3 displays the evolution of the 2σ lower limit as a function of the CIB uncertainty scaling factor. A scaling factor x means that the errors are assumed to be x times smaller than currently estimated. The lower bound is, as expected, improved by smaller uncertainties and reaches a limit when the propagation of errors induced by the CIB uncertainties becomes much smaller than the effect of gamma-ray uncertainties associated with the measurement of the AGN spectrum. For each point of the plot, 1000 Monte-Carlo realizations have been computed and analyzed according to the method described in the previous section.

In the near future, measurements should indeed be improved between 60 and 110 μm thanks to *Herschel* (see, *e.g.* Franceschini *et al.* (2006)) and, in the far future, between 5 and 60 μm thanks to *JWST* (see, *e.g.* Windhorst *et al.* (2006)). This could make this approach quite competitive.

Finally, if gamma telescopes become so efficient that many more blazars are detected, each of them exhibiting a redshift-dependant cutoff in the spectrum, then the approach suggested by Salamon *et al.* (1994) could be tractable and would lead not only to a bound but to a measurement of the Hubble parameter. However, to distinguish between a CIB absorption effect and an intrinsic cutoff, a very large number of data points would be required.

REFERENCES

- Aharonian, F.A. *et al.*, 1997, A&A 327, L5
 Aharonian, F.A. *et al.*, 1999a, A&A 342, 69
 Aharonian, F.A., *et al.*, 1999b, A&A, 349, 11
 Aharonian, F.A. *et al.*, 2000, New Astron., 5, 377
 Aharonian, F.A. *et al.*, 2005, A&A, 430, 865
 Aharonian, F.A. *et al.*, 2005, A&A, 441, 465
 Aharonian, F.A. *et al.*, 2006, Nature, 440, 1018
 Aharonian, F.A. *et al.*, 2007, A&A, 470, 475
 Aharonian, F.A. *et al.*, 2007, ApJ, 664, L71
 Bernstein, R.A., Freedman, W.L. & Madore, B.F., 2002, ApJ, 571, 107
 Bastieri, D., 2008, Nucl. Instrum. Meth. A 588, 1
 Biller, S.D. *et al.*, 1998, Phys. Rev. Lett., 80, 2992
 Blanchard, A., Douspis, M., Rowan-Robinson, M., Sarkar, S., 2003, A&A, 412, 35
 Bonamente, M., Joy, M.K., Laroque, S.J., Carlstrom, J.E., Reese, E.D., Dawson, K.S., 2006, ApJ, 647, 25
 Böttcher, M., Basu, S., Joshi, M., Villata, M., 2007, ApJ, 670, 968
 Cambresy, L., Reach, W.T., Beichman, C.A. & Jarett, T.H., 2001, ApJ, 555, 563
 Catanese, M. *et al.*, 1997, ApJ, 487, L143
 Chiaberge, M., Capetti, A. & Celloti, A., 2001, MNRAS, 318, 493
 Coppi, S. & Aharonian F.A., 1999, ApJ 521, L33
 Djannati-Ataï, A. *et al.*, 1999, A&A, 350, 17
 Dole, H. *et al.*, 2004, ApJS, 154, 87
 Dole, H. *et al.*, 2006, A&A, 451, 2, 417
 Elbaz, D. *et al.*, 1999, A&A, 351, L37
 Freedman, W.L. *et al.*, 2001, ApJ 553, 47
 Franceschini, A., Vaccari, M., Berta, S., Rodighiero, G., & Lonsdale, C., 2006, Proceeding of the conference "Studying Galaxy Evolution with Spitzer and Herschel", Crete, Eds. V. Charmandaris, D. Rigopoulou, N. Kylafis [arXiv:astro-ph/0610897v1]
 Ghisellini, G., Foschini, L., Tavecchio, F. & Pian, E., 2007, MNRAS, 382, L82
 Gispert, R., Lagache, G. & Puget, J.L., 2000, A&A 360, 1
 Giommi, P., Capalbi, M., Fiocchi, M. *et al.*, 2002, in Blazar Astrophysics with BeppoSax and Other Observations, ed. P. Giommi, E. Massaro & G. Palumbo
 Gorjian, V., Wright, E.L., Chary, R.R., 2000, ApJ, 536, 550
 Guy, J., Renault, C., Aharonian, F., Rivoal, M., Tavernet, J.-P., 2000, A&A 359, 419
 Hauser, M.G. *et al.*, 1998, ApJ 508, 25
 Hauser, M.G. & Dwek, E., 2001 ARA&A, 37, 249
 Hayashida, N. *et al.*, 1998, ApJ 504, 71
 Heitler, W., 1960, The Quantum Theory of Radiation, (Oxford)
 Jackson, N., 2007, Living Rev. Rel. 10, 4
 Jones, M.E. *et al.*, 2005, MNRAS, 357, 518
 Kashlinsky, A., 2005, Phys. Rept., 409, 361
 Komatsu, E. *et al.*, 2008, submitted to ApJ [arXiv:0803.0547]
 Konopelko, A.K., Kirk, J.G., Stecker, F.W. & Mastichiadis, A., 1999, ApJ 518, L13
 Krennrich, F., 2007, J. Phys. Conf. Ser., 60, 34
 Lagache, G., Haffner, L.M., Reynolds, R.J., Tufte S.L., 1999, A&A, 354, L247
 Lamer, G. & Wagner, S.J., 1998, A&A 331, L13
 Latronico, L., Spandre, G., 2007, Nucl. Instrum. Meth. A, 581, 160
 Lenain, J.-P., Boisson, C., Sol, H. & Katarzynski, 2008, A&A, 478, 111
 Matilla, K., 2003, ApJ, 591, 119
 Nikishov, A. I. 1962, Soviet Physics JEPT 14, 2
 Nugent, P., Branch, D., Baron, E., Fisher, A., Vaughan, T., Hauschildt, P.P., 1995, Phys. Rev. Lett., 75, 394

- Oguri, M., 2007, ApJ, 660, 1
Papovich, C. *et al.*, 2004, ApJS, 154, 70
Pian, E. *et al.*, 1998, ApJ 492, L17
Puget, J.L., Abergel, A., Bernard, J.P., Boulanger, F., Burton, W.B., Desert, F.X., Hartmann, D., 1996 A&A; 308, L5
Rieke, G.H., Young, E.T., Engelbracht, C.W., 2004, AJPS, 154, 25
Riess, A.G. *et al.*, 2005, ApJ, 627, 579
Renault, C., Barrau, A., Lagache, G. & Puget, J.L., 2001, A & A, 371, 771
Saha, A., Coles, J., Macció, A.V. & Williams, L.L.R., ApJL, 650, L17
Samuelson, F. W., Biller S. D., Bond I. H., Boyle P.J., 1998, ApJ, 501, L17
Sandage, A. *et al.*, 2006, ApJ, 653, 843
Saugé, L., PhD thesis from Grenoble Unive., 2004
Salamon, M.H., Stecker, F.W., de Jager, O.C. 1994, ApJL, 423, L1
Stanev, T., Franceschini, A., 1998, ApJ 494, L159
Stecker, F.W. & De Jager, O.C., 1998, A&A, 334, L85
Stecker, F.W. & De Jager, O.C., & Salamon 1992, ApJ, 390, L49
Stecker, F.W., Baring, M.G., Summerlin, E.J., 2007, ApJ, 667, L29
Tegmark, M. *et al.*, 2006, Phys. Rev. D, 74, 123507
Türler, M. *et al.*, 1999, A&AS, 134, 89
Werner, M.W., *et al.*, 2004, ApJS, 154, 1
Windhorst, R.A., Cohen, S.H., Jansen, R.A., Conselice, C. & Yan, H.-J., 2006, New Astron. Rev., 50, 113
Wright, E.L., 2001, ApJ, 553, 748